
The inactive X chromosome as a female protector in autism and beyond

In the format provided by the authors and unedited

Supplementary Note

Underdiagnosis of autistic females and increased male susceptibility to X-linked causes of autism are frequently invoked to explain the male bias of autism. Here, we present evidence that these factors cannot fully account for autism's male bias.

Underdiagnosis of autistic females

Females often face greater barriers to receiving an autism diagnosis compared to males, due to factors such as clinician bias, gendered behavioral expectations, and sex differences in the extent to which core symptoms of autism are expressed versus “masked”¹. This has led some to speculate that the male preponderance of autism is driven by underdiagnosis of autistic females². While diagnostic bias almost certainly contributes to the male skew of autism, a male-biased sex ratio persists even when only considering autism cases that are readily clinically ascertainable and thus less susceptible to significant diagnostic bias. For example, the Baby Siblings Research Consortium screened for autism in the younger siblings of over one thousand autistic individuals and still found a 3M:1F sex bias among affected siblings^{3,4}.

Bias is also less likely to influence the diagnosis of profoundly affected autistic individuals, such as those with intellectual disability (ID). Consistently, sex ratio estimates in autism with ID range from 2– 3.5M:1F, whereas sex ratio estimates for autism without ID are higher, from 4-9M:1F^{1,5-8}. It is tempting to speculate that this increased sex ratio in less profoundly affected individuals reflects diagnostic bias, but it is important to note that there are also significant differences in the genetic architecture of autism with and without ID that could influence a sex bias⁹⁻¹². For example, autism without co-occurring intellectual disability (ID) has been more strongly linked to common polygenic variation, whereas autism with ID is more often

associated with damaging, frequently *de novo* mutations in highly constrained genes^{9,13}.

Regardless, it is clear that even readily ascertainable cases of autism show a significantly male-biased sex ratio.

Finally, the male bias of autism is consistently observed across countries with diverse cultures and gender norms. A recently published study of the global prevalence of autism reported a male bias in each of over two hundred surveyed countries and regions from the Americas, Europe, Asia, Africa, and Australasia¹⁴. Taken together, these findings suggest that sociocultural factors or underdiagnosis of autistic females cannot fully account for the male bias in autism.

Enhanced male vulnerability to X-linked causes of autism

Another common explanation for autism's male bias is the increased vulnerability of males to X-linked – and particularly X-linked recessive – causes of autism, as males have just one copy of X-linked genes. Indeed, 33 of 91 X-linked autism genes are reported to follow a strictly recessive mode of inheritance (MOI) in OMIM¹⁵, suggesting a substantial contribution of X-linked recessive mutations to autism (**Fig. S1A, Table S1, and Supplementary Methods**). However, even X-linked recessive (XLR) mutations can sometimes affect females due to variable expressivity, or to skewing of X-chromosome inactivation (XCI). For example, a detailed phenotypic analysis of 19 females heterozygous for mutations in *KDM5C*, an X-linked gene classically thought to cause autism in a recessive manner¹⁶, found that 15 of the 19 exhibited moderate ID¹⁷. Thus, while XLR mutations certainly contribute to male risk for autism, they also can increase risk for autism and related neurodevelopmental symptoms in females.

Moreover, 13 of the 91 X-linked autism genes (**Fig. S1A**) exhibit an X-linked dominant (XLD) MOI. For several reasons, these XLD forms would be expected to bias autism toward females. First, many XLD causes of autism in females are typically lethal in males. For example, protein-truncating variants in *MECP2* and *DDX3X* cause syndromic autism in females (Rett and *DDX3X* syndrome, respectively) but are nearly always fatal in males^{18,19}. Second, paternal X chromosomes, which are transmitted only to daughters, tend to carry more *de novo* mutations than maternal X chromosomes, which are transmitted to both sons and daughters²⁰. Therefore, females are more likely than males to inherit a *de novo* mutation in an X-linked autism gene with a dominant MOI²¹. Consequently, even XLD causes of autism that are not lethal in males will disproportionately affect females. This female-biased prevalence has been consistently reported in the literature and acts in opposition to the observed male bias in autism. For example, four sex-stratified analyses of *de novo* mutations (DNMs) in X-linked genes associated with autism or developmental disorders (DDs) identified more female-enriched genes (genes significantly enriched for DNMs in affected females only) than male-enriched genes, despite males constituting the majority of the cohort in each of the four studies (**Table S2**)²¹⁻²⁴.


Finally, 27 of the 91 X-linked autism genes follow neither a strictly recessive nor dominant MOI (X-linked MOI; **Fig. S1A**). As a result, mutations in these genes are not expected to uniformly contribute to a male bias in autism^{21,25,26}. Considering all of these observations together, it appears unlikely that enhanced male vulnerability to X-linked mutations can explain the strong male bias of autism.

This conclusion is further supported by mathematical modeling of autism's male bias. Previous studies of general DDs and ID, both with male-biased sex ratios of 1.4M:1F, have shown that X-linked mutations cannot account for the excess of male cases in these


conditions^{21,27}. Using the approach described by Ropers and Hamel²⁷, we find that 67 – 75% of male cases of autism would require X-linked etiologies that do not affect females to explain a sex ratio of 3-4M:1F (**Fig. S1B** and **Supplementary Methods**). This proportion is much higher than what has been reported in previous studies. For example, the most common X-linked cause of autism in males, Fragile X Syndrome, accounts for only 1 – 6% of all cases of autism²⁸. Broader searches for X-linked autism genes have been of great interest but have largely failed to identify loci that explain a substantial proportion of male autism risk. For example, one of the largest autism exome sequencing studies to date ($n = 11,986$ autism cases; 17% female) found no X-linked gene with a significant excess of deleterious variants in sex-stratified case-control analyses²⁹. In contrast, 102 significant autism risk genes were identified on the autosomes. Subsequent autism exome sequencing studies have yielded similar results^{30,31}. Even targeted searches for X-linked variants that disproportionately affect males have only explained an additional 2-3% of male cases of autism³². Thus, though some male autism cases are attributable to X-linked etiologies, and efforts to identify additional X-linked risk factors are ongoing^{33,34}, it seems implausible that enough X-linked variation remains undiscovered to explain the observed male bias of autism.

Supplementary Figures

A



B



Supplementary Figure 1. Enhanced male susceptibility to X-linked, and particularly X-linked recessive, causes of autism cannot explain its male bias. (A) Pie chart of modes of inheritance (MOIs) of X-linked autism genes. Most X-linked autism genes do not show a clear X-linked recessive mode of inheritance. X-linked autism genes obtained from *SFARI Gene*³⁵.

MOIs obtained from *OMIM*¹⁵. X-linked dom: X-linked dominant. **(B)** Mathematical modeling²⁷ of theoretical proportion of male autism cases required to be attributable to X-linked mutations that do not affect females to explain the reported 3-4M:F sex ratio of autism. 67 - 75% of male cases of autism would need to be due to X-linked causes to fully explain this male bias, a proportion far above what has been seen in sequencing studies of autistic males.

Supplementary Methods

Analysis of modes of inheritance of X-linked autism genes

A list of 91 X-linked genes associated with autism was obtained from the Q4 2024 release of the *SFARI Gene*³⁵ database. *OMIM* (omim.org)¹⁵ was used to systematically identify their mode of inheritance (MOI). 18 X-linked genes (*AGTR2*, *CDK16*, *CD99L2*, *DDX53*, *FAM47A*, *IL1RAPL2*, *MAGEC3*, *MAOB*, *PCDH11X*, *PJAI*, *PLXNA3*, *PTCHD1-AS*, *RHOXF1*, *SLC7A3*, *SYAP1*, *TRPC5*, *TSPYL2*, *VSIG4*) had no associated OMIM phenotypes. For all other X-linked genes with at least one associated OMIM phenotype, genes were grouped into the following categories based on OMIM reported MOIs: (1) X-linked recessive only (XLR), (2) X-linked dominant (XLD), (3) multiple modes of inheritance reported (*i.e.*, both XLR and XLD), and (4) X-linked only (*i.e.*, only X-linked reported in OMIM, with no reporting of either XLD or XLR MOIs).

Modeling the contribution of X-linked mutations needed to explain the observed male bias of autism

As demonstrated by Ropers and Hamel²⁷, the proportion of males that must have an X-linked mutation that does not affect females to fully explain the male case excess in a condition with a male-to-female ratio of r is given by $(r-1)/r$. Under the assumption of a male-to-female ratio of 3-4:1 in autism, this suggests that 67-75% of autistic males would have to have an X-linked cause, respectively, to fully explain the observed male bias.

References

1. Werling DM. The role of sex-differential biology in risk for autism spectrum disorder. *Biol. Sex Differ.* **7**, 58 (2016).
2. Furfaro H. The female protective effect, explained. *The Transmitter: Neuroscience News and Perspectives* (2019). Available at: <https://www.thetransmitter.org/spectrum/female-protective-effect-explained/>
3. Ozonoff S et al. Recurrence risk for autism spectrum disorders: a Baby Siblings Research Consortium study. *Pediatrics* **128**, e488–e495 (2011).
4. Messinger DS et al. Early sex differences are not autism-specific: a Baby Siblings Research Consortium (BSRC) study. *Mol. Autism* **6**, 32 (2015).
5. Loomes R, Hull L and Mandy WPL. What is the male-to-female ratio in autism spectrum disorder? A systematic review and meta-analysis. *J. Am. Acad. Child Adolesc. Psychiatry* **56**, 466–474 (2017).
6. Lord C and Schopler E. Brief report: differences in sex ratios in autism as a function of measured intelligence. *J. Autism Dev. Disord.* **15**, 185–193 (1985).
7. Volkmar FR, Szatmari P and Sparrow SS. Sex differences in pervasive developmental disorders. *J. Autism Dev. Disord.* **23**, 579–591 (1993).
8. Yeargin-Allsopp M et al. Prevalence of autism in a US metropolitan area. *JAMA* **289**, 49–55 (2003).
9. Warrier V et al. Genetic correlates of phenotypic heterogeneity in autism. *Nat. Genet.* **54**, 1293–1304 (2022).
10. Bulik-Sullivan B et al. An atlas of genetic correlations across human diseases and traits. *Nat. Genet.* **47**, 1236–1241 (2015).

11. Clarke T-K et al. Common polygenic risk for autism spectrum disorder is associated with cognitive ability in the general population. *Mol. Psychiatry* **21**, 419–425 (2016).
12. Hagenaars SP et al. Shared genetic aetiology between cognitive functions and physical and mental health in UK Biobank (N = 112,151) and 24 GWAS consortia. *Mol. Psychiatry* **21**, 1624–1632 (2016).
13. Grove J et al. Identification of common genetic risk variants for autism spectrum disorder. *Nat. Genet.* **51**, 431–444 (2019).
14. Santomauro DF et al. The global epidemiology and health burden of the autism spectrum: findings from the Global Burden of Disease Study 2021. *Lancet Psychiatry* **12**, 111–121 (2025).
15. Hamosh A, Scott AF, Amberger JS, Bocchini CA and McKusick VA. Online Mendelian Inheritance in Man (OMIM), a knowledgebase of human genes and genetic disorders. *Nucleic Acids Res.* **33**, D514–D517 (2005).
16. Schenkel LC et al. Peripheral blood epi-signature of Claes-Jensen syndrome enables sensitive and specific identification of patients and healthy carriers with pathogenic mutations in KDM5C. *Clin. Epigenetics* **10**, 21 (2018).
17. Carmignac V et al. Further delineation of the female phenotype with KDM5C disease causing variants: 19 new individuals and review of the literature. *Clin. Genet.* **98**, 43–55 (2020).
18. Gold WA et al. Rett syndrome. *Nat. Rev. Dis. Primers* **10**, 1–17 (2024).
19. Kennis MGP et al. DDX3X-related neurodevelopmental disorder in males – presenting a new cohort of 19 males and a literature review. *Eur. J. Hum. Genet.* **33**, 980–988 (2025).

20. Pinto N et al. Paternal and maternal mutations in X-STRs: a GHEP-ISFG collaborative study. *Forensic Sci. Int. Genet.* **46**, 102258 (2020).
21. Martin HC et al. The contribution of X-linked coding variation to severe developmental disorders. *Nat. Commun.* **12**, 627 (2021).
22. Zhang Y et al. Genetic evidence of gender difference in autism spectrum disorder supports the female-protective effect. *Transl. Psychiatry* **10**, 11 (2020).
23. Turner TN et al. Sex-based analysis of de novo variants in neurodevelopmental disorders. *Am. J. Hum. Genet.* **105**, 1274–1285 (2019).
24. Wang T et al. Integrated gene analyses of de novo variants from 46,612 trios with autism and developmental disorders. *Proc. Natl. Acad. Sci. U. S. A.* **119**, e2203491119 (2022).
25. Dobyns WB et al. Inheritance of most X-linked traits is not dominant or recessive, just X-linked. *Am. J. Med. Genet. A* **129A**, 136–143 (2004).
26. Basava S, Billington CJ Jr, Carrel L, Biesecker LG and Dobyns WB. Patterns of X-linked inheritance: a new approach for the genome era. *Genet. Med.* **27**, 101384 (2025).
27. Ropers H-H and Hamel BCJ. X-linked mental retardation. *Nat. Rev. Genet.* **6**, 46–57 (2005).
28. Kaufmann WE et al. Autism spectrum disorder in fragile X syndrome: cooccurring conditions and current treatment. *Pediatrics* **139**, S194–S206 (2017).
29. Satterstrom FK et al. Large-scale exome sequencing study implicates both developmental and functional changes in the neurobiology of autism. *Cell* **180**, 568–584.e23 (2020).
30. Zhou X et al. Integrating de novo and inherited variants in 42,607 autism cases identifies mutations in new moderate-risk genes. *Nat. Genet.* **54**, 1305–1319 (2022).

31. Fu JM et al. Rare coding variation provides insight into the genetic architecture and phenotypic context of autism. *Nat. Genet.* **54**, 1320–1331 (2022).
32. Wang S et al. Rare X-linked variants carry predominantly male risk in autism, Tourette syndrome, and ADHD. *Nat. Commun.* **14**, 8077 (2023).
33. Scala M et al. Genetic variants in DDX53 contribute to autism spectrum disorder associated with the Xp22.11 locus. *Am. J. Hum. Genet.* **112**, 154–167 (2025).
34. Mendes M et al. Chromosome X-wide common variant association study in autism spectrum disorder. *Am. J. Hum. Genet.* **112**, 135–153 (2025).
35. Abrahams BS et al. SFARI Gene 2.0: a community-driven knowledgebase for the autism spectrum disorders. *Mol. Autism* **4**, 36 (2013).